



Application of computational systems biology to explore environmental toxicity hazards.

Audouze, Karine Marie Laure; Grandjean, Philippe

Published in:
Environmental Health Perspectives

Link to article, DOI:
[10.1289/ehp.1103533](https://doi.org/10.1289/ehp.1103533)

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Audouze, K. M. L., & Grandjean, P. (2011). Application of computational systems biology to explore environmental toxicity hazards. *Environmental Health Perspectives*, 119(12), 1754-1759.
<https://doi.org/10.1289/ehp.1103533>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Application of Computational Systems Biology to Explore Environmental Toxicity Hazards

Karine Audouze¹ and Philippe Grandjean^{2,3}

¹Center for Biological Sequence Analysis, Department of Systems Biology, Technical University of Denmark, Lyngby, Denmark;

²Institute of Public Health, University of Southern Denmark, Odense, Denmark; ³Department of Environmental Health, Harvard School of Public Health, Boston, Massachusetts, USA

BACKGROUND: Computer-based modeling is part of a new approach to predictive toxicology.

OBJECTIVES: We investigated the usefulness of an integrated computational systems biology approach in a case study involving the isomers and metabolites of the pesticide dichlorodiphenyltrichloroethane (DDT) to ascertain their possible links to relevant adverse effects.

METHODS: We extracted chemical–protein association networks for each DDT isomer and its metabolites using ChemProt, a disease chemical biology database that includes both binding and gene expression data, and we explored protein–protein interactions using a human interactome network. To identify associated dysfunctions and diseases, we integrated protein–disease annotations into the protein complexes using the Online Mendelian Inheritance in Man database and the Comparative Toxicogenomics Database.

RESULTS: We found 175 human proteins linked to *p,p'*-DDT, and 187 to *o,p'*-DDT. Dichlorodiphenyldichloroethylene (*p,p'*-DDE) was the metabolite with the highest number of links, with 52. We grouped proteins for each compound based on their disease annotations. Although the two data sources differed in linkage to diseases, integrated results predicted that most diseases were linked to the two DDT isomers. Asthma was uniquely linked with *p,p'*-DDT, and autism with *o,p'*-DDT. Several reproductive and neurobehavioral outcomes and cancer types were linked to all three compounds.

CONCLUSIONS: Computer-based modeling relies on available information. Although differences in linkages to proteins may be due to incomplete data, our results appear meaningful and suggest that the parent DDT compounds may be responsible for more disease connections than the metabolites. The findings illustrate the potential use of computational approaches to toxicology.

KEY WORDS: computational biology, DDT, genomics, proteomics, systems biology. *Environ Health Perspect* 119:1754–1759 (2011). <http://dx.doi.org/10.1289/ehp.1103533> [Online 17 August 2011]

In its report, *Toxicity Testing in the 21st Century*, the National Research Council called for the development of new approaches to human health risk assessment that would rely, in part, on computer-based models rather than animal testing and epidemiology (National Research Council 2007). Although these recommendations were timely and visionary, progress has been slow, possibly because of the need for elaborate validation of models before the adoption of a new approach to predictive toxicology. However, toxicological databases and computational methods have developed further and now seem to be even better suited for applications in environmental health research. Through international efforts, publicly available databases have been combined and fine-tuned to provide linked information regarding chemical names, synonyms, chemical structures, hazards, chemical exposures, and potential risks to human health within several categories: acute, developmental toxicity, reproductive toxicity, and cancer (Judson et al. 2008). As a complement to these databases, ChemProt, a new disease chemical biology database (Center for Biological Sequence Analysis 2011), provides information on chemical links to proteins along with chemical names, chemical structures, and diseases

(Taboureau et al. 2011). These data resources can be used to develop computational models for predicting toxicological end points or possible biological mechanisms.

Various models already have been developed, most of them structure-based. Chemicals are grouped according to their similar structural features or fragments that can be related to a particular toxicity end point, for example, ToxMatch (Pavan and Worth 2008) and Derek Nexus (Lhasa Limited 2011), or by using quantitative structure–activity relationships approaches, as in Computer Assisted Evaluation of Industrial Chemical Substances According to Regulations (CAESAR) (Cassano et al. 2010). In parallel, systems chemical biology has emerged as a field that integrates chemical information with biological databases (Oprea et al. 2007). It has therefore become possible to generate a generic computational systems biology model that aims at revealing the underlying molecular mechanisms of xenobiotics and the biological pathways they may disrupt (Audouze et al. 2010). In this approach, toxicogenomics data are combined with systems biology information to provide a high-confidence human protein–protein association network. We previously used this integrative systems biology method

to decipher unexpected relationships between di(2-ethylhexyl)phthalate and gamma-aminobutyric acid receptors (Audouze et al. 2010). Similarly, Gohlke et al. (2009) developed a network of complex diseases to integrate molecular pathways associated with both genetic and environmental factors. However, these computational predictions and others like them must be interpreted in light of the caveat that the databases used contain available information only and that metabolism and other factors may affect the chemical–protein interactions. Given the new developments and our promising initial experience, we have attempted to illustrate the current potential of advanced computational systems biology to assess the potential hazards associated with a group of environmental chemicals with substantial, although incomplete, toxicological and epidemiological information.

The pesticide dichlorodiphenyltrichloroethane (DDT) was marketed as an ideal insecticide with long-term protection; however, the environmental persistence of DDT and its metabolite dichlorodiphenyldichloroethylene (DDE) resulted in bioaccumulation and delayed or latent adverse effects. Experimental studies have explored mainly the effects of DDT dosages, whereas epidemiology research primarily has used the serum concentration of the DDE metabolite as an exposure biomarker. The DDE concentration may reflect, to some degree, past exposures to the parent DDT compound that had been metabolized later, but it may also originate directly from DDE residues in food. Thus, concentrations present at the time of blood sampling may represent neither the correct chemical species nor the active dose present at the time when a possible adverse effect was initiated. This conundrum is difficult to resolve from epidemiology

Address correspondence to K. Audouze, Center for Biological Sequence Analysis, Department of Systems Biology, Technical University of Denmark, DK-2800 Lyngby, Denmark. Telephone: 45 45252472. Fax: 45 45931585. E-mail: karine@cbs.dtu.dk

Supplemental Material is available online (<http://dx.doi.org/10.1289/ehp.1103533>).

This work was supported by the Villum Kann Rasmussen Foundation, the EU Innovative Medicines Initiative Joint Undertaking (eTOX), and the National Institute for Environmental Health Sciences (grants ES 012199 and ES 013692).

The authors declare they have no actual or potential competing financial interests.

Received 7 February 2011; accepted 8 August 2011.

data alone, and the experimental data do not cover all the potential outcomes for the relevant isomers and metabolites [Agency for Toxic Substances and Disease Registry (ATSDR) 2002]. We therefore chose to carry out computational chemistry analyses of the DDT compounds from a human systems biology perspective. Our dual purpose was to obtain new information that might link relevant outcomes to specific DDT compounds and to ascertain to what degree the currently available data sources would allow such analyses using a systems chemical biology approach. Although this study was not intended as a formal validation study, we consider a family of compounds for which extensive although somewhat equivocal epidemiological and experimental evidence exists, with the result that our analysis includes suspected causal agents and anticipated negative controls.

Methods

Using data from the ToxProfile of the ATSDR (2002) with the 2008 appendix and literature searches of PubMed, we chose to concentrate on inflammatory, reproductive and endocrine, neurobehavioral, and malignant diseases. DDT isomers and metabolites were examined by multi-step data integration in the systems chemical biology approach (Figure 1). In short, in the first step, we extracted existing knowledge from a disease chemical biology database to generate compound-specific human protein networks. In the second step, protein enrichment, we used a high-confidence set of experimental human protein–protein interactions to identify protein complexes. In the final step, we ranked the diseases predicted to be linked to the DDT compounds using an integration of protein–disease annotations in the protein complexes.

More specifically, first we extracted the human protein–chemical associations from the newly established disease chemical biology database, ChemProt (Taboureau et al. 2011), a compilation of multiple chemical–protein annotation resources that contains 700,000 unique chemicals connected with 30,578 proteins. This database assembles chemical–protein connections from multiple sources, such as ChEMBL (de Matos et al. 2010), BindingDB (Liu et al. 2007), PDSP Ki database (Roth et al. 2004), and PubChem bioassays (Wheeler et al. 2007). The integrated associations comprise both binding and separate gene expression data, as the deregulation of a gene by a chemical may not be due necessarily to a physical interaction between the compound and the protein, e.g., in the form of binding, but could entail a response at a cellular level. ChemProt can predict uncharacterized chemical–protein interactions based on the similar structure of compound pairs. In the present study, only chemical–protein

relationships with experimental support were kept for the analysis, while otherwise unsupported, predicted connections were disregarded.

Second, we explored protein–protein interactions through an in-house human interactome network based on experimental data from humans and 21 model organisms (Lage et al. 2007, 2008). Using a probabilistic confidence scoring scheme, all interactions in the human interactome have been validated against a gold standard (Rual et al. 2005). We used an updated version of the protein–protein interaction network consisting of refined experimental proteomics data (Lage et al. 2010). The current interactome contains 507,142 unique protein–protein interactions (PPIs). These data are derived from sources such as the Biomolecular Interaction Network Database (BIND; Bader et al. 2003), Biological General Repository for Interaction Datasets (BioGRID; Stark et al. 2006), Molecular Interaction (MINT) database (Zanzoni et al. 2002), DIP_FULL dataset (Salwinski et al. 2004), Human Protein Reference Database (HPRD; Mishra et al. 2006), IntAct (an open source molecular interaction database) (Hermjakob et al. 2004), Mammalian Protein–Protein Interaction (MPPI) database (Pagel et al. 2005), MPact [the Munich Information Center for Protein Sequences (MIPS) protein interaction resource on yeast] (Guldener et al. 2006), Reactome (Joshi-Tope et al. 2005), and Kyoto Encyclopedia of Genes and Genomes (KEGG) databases (Kanehisa et al. 2006). Data are transferred between organisms by using the InParanoid Eukaryotic Ortholog Groups' database (InParanoid 2009; O'Brien et al. 2005). Among the PPIs included, 414,543 are listed at least once as interactions between two human proteins. The human interactome includes 22,997 genes. Among the gene products, 14,441 human proteins have known interactions with other proteins; the others are singletons. In addition to direct interactions between known proteins, we also included indirect interactions by considering proteins involved in common pathways. This step used a statistical procedure shown to provide optimal network significance (Lage et al. 2010). As a result, the list of relevant proteins associated with each compound is extended by inclusion of their known first-order protein–protein interaction partners and other proteins participating in the same pathways.

Third, to identify diseases or dysfunctions associated with the specific chemicals examined (i.e., *p,p'*-DDT, *o,p'*-DDT, and *p,p'*-DDE), we integrated protein-specific information related to their involvement in disease pathogenesis with the proteins identified. The Online Mendelian Inheritance in Man (OMIM) database [Hamosh et al. 2005; National Center for Biotechnology Information (NCBI) 2011b] and the Comparative Toxicogenomics

Database (CTD 2011; Davis et al. 2011) were selected as the most useful sources of protein–disease information. The OMIM database is a highly reliable compilation of genetic variants from medical and genetics publications. The CTD database contains both direct (manually curated) and inferred gene–disease relationships and may therefore provide less certain




Figure 1. Overview of the systems chemical biology three-step approach. (1) Extraction of existing knowledge using a disease chemical biology database (ChemProt) to generate chemical–protein networks for *p,p'*-DDT. (2) Creation of protein complexes by protein enrichment using a high-confidence set of experimental protein–protein interactions. (3) Statistical ranking of diseases (D) known or predicted to be linked to *p,p'*-DDT after integration of protein–disease annotations to protein complexes based on information in the Online Mendelian Inheritance in Man (OMIM) database and the Comparative Toxicogenomics Database (CTD).

associations. Whereas direct associations derive from experimental models or epidemiological studies, inferred relationships are established through indirect evidence. Thus, if gene A is associated with disease B, and gene A has a curated interaction with chemical C, then chemical C has a curated relationship with disease B (Davis et al. 2011). These inferred relationships are more explorative and allow hypothesis generation using protein–disease connections in the chemical space.

The different strengths of the two databases, OMIM and CTD, are illustrated by comparison of the data sets. The OMIM (version 2009) contains 3,748 connections involving 2,728 unique diseases, while the CTD (version 2009) contains 252,056 (hypothetical) protein–disease associations involving 2,580 unique diseases. We translated diseases having a Medical Subject Heading identification (MeSH ID) number in CTD into OMIM ID numbers to the extent possible, given that a single MeSH ID number may correspond to several OMIM ID numbers. Based on OMIM ID numbers, 1,934 diseases are shared between the two databases, 2,042 proteins are common to both, and they correspond to 2,528 shared protein–disease associations of 255,804 interactions listed. Because of the limited overlap, we used both resources to explore the predictions in terms of protein–disease relationships and biological confidence with regard to grouping proteins within a binary metric distance (Audouze et al. 2010).

Using R software (R Development Core Team 2010), *p*-values were calculated for each group of proteins, assuming a hypergeometric distribution. To take into account the large numbers of potential protein–disease connections in the CTD, we performed Bonferroni adjustments of all *p*-values for protein clusters based on the CTD. For this source of information, we chose a classical family-wise error rate of 0.05 as cutoff to adjust *p*-values. We mapped all human proteins identified to EntrezGene

(NCBI 2011a) identifiers using the Clone/ GeneID converter (Alibes et al. 2007).

Results

Using the ChemProt database, we extracted 38 relevant human proteins for *p,p'*-DDT, 83 for *o,p'*-DDT, and 18 for *p,p'*-DDE [for details, see Supplemental Material, Table 1 (<http://dx.doi.org/10.1289/ehp.1103533>)]. We used the resulting three lists of proteins independently to create three human protein networks. For *o,p'*-DDE, only 11 proteins were identified, and all had already been identified earlier as targets for *p,p'*-DDE. Likewise, *p,p'*-DDD was connected with only 10 proteins, and *o,p'*-DDD with only three [progesterone receptor, androgen receptor, and estrogen receptor 1, the latter two overlapping with *p,p'*-DDD]. No information was obtained on the methylsulfonyl metabolites. Our data analysis therefore concentrated on the three substances first mentioned.

We generated a human protein network for each chemical by determining protein–protein interaction partners associated with each protein network. Proteins with a GeneEntrez ID were retained. We identified 175 proteins for the *p,p'*-DDT analysis, 187 proteins for *o,p'*-DDT, and 52 proteins for *p,p'*-DDE (Table 1). Disease enrichment based on OMIM and CTD protein–disease annotations led to identification of diseases associated with each chemical (Table 2). Some diseases were not significantly associated with a chemical (e.g., when linked only via one protein). As anticipated, the CTD provided many more potential disease associations than the OMIM. This difference between the OMIM- and CTD-based predictions likely reflects the extent of uncertain and incomplete evidence within the two data sources. The specific and

overlapping disease annotations for *p,p'*-DDT and *p,p'*-DDE within disease clusters for the two different data sources are shown in Figure 2. Most of the diseases predicted for *o,p'*-DDT overlapped with predictions for *p,p'*-DDT. Overall, 35 diseases (based on OMIM data) and 210 diseases (based on CTD data) appeared to be unique for *p,p'*-DDT. All OMIM-based predictions had higher *p*-values than CTD-based predictions, but these differences should be interpreted in light of the fewer known proteins in the OMIM database associated with the diseases predicted. For example, asthma was connected with *p,p'*-DDT via a single protein, tumor necrosis factor, in the OMIM database (*p* = 0.147), whereas CTD information predicted asthma via connections through 48 proteins (*p* = 0.002) [for details, see Supplemental Material, Table 2 (<http://dx.doi.org/10.1289/ehp.1103533>)]. In contrast, only a few diseases appeared to be unique for *p,p'*-DDE: 1 based on CTD data (i.e., coronary heart disease), and 8 based on OMIM data.

To synthesize this information, we focused on the four major categories of disease phenotypes previously linked to DDT exposures: inflammatory, reproductive and endocrine, neurobehavioral, and carcinogenic (Figure 3). [See Supplemental Material, Table 2 (<http://dx.doi.org/10.1289/ehp.1103533>) for details on the *p*-values (adjusted when based on CTD data), sources of protein–disease information (CTD or OMIM), and specific genes linked to the proteins.] Depending on the source of protein–disease annotations, the phenotypes may be more or less specific, but all annotations identified in the databases were kept to avoid any subjective judgment in data extraction. As expected, the predictions varied somewhat between the two databases, OMIM and CTD.

Table 1. Number of proteins associated with *p,p'*-DDT, *o,p'*-DDT, and *p,p'*-DDE within the different steps of the systems biology procedure.

Chemical name	Number of proteins		
	ChemProt	Interactome (PPIs)	Mapped to EntrezGene
<i>p,p'</i> -DDT	38	182 (381)	175
<i>o,p'</i> -DDT	83	189 (235)	187
<i>p,p'</i> -DDE	18	56 (92)	52

Table 2. Number of diseases associated with *p,p'*-DDT, *o,p'*-DDT, and *p,p'*-DDE using the OMIM and the CTD databases.

Chemical name	OMIM	CTD
<i>p,p'</i> -DDT	50	271
<i>o,p'</i> -DDT	45	77
<i>p,p'</i> -DDE	25	62




Figure 2. Venn diagram showing the number of diseases overlapping between *p,p'*-DDE and *p,p'*-DDT, using disease annotations extracted from the OMIM and the CTD databases. Data on *o,p'*-DDT are not shown, as most of its disease links overlapped with *p,p'*-DDT.

Overall, of 175 proteins linked to *p,p'*-DDT after exploration of the human interactome in the CTD, only 29 were retrieved using the OMIM database. Among those 29 proteins, one was associated with asthma among the total of 13 connected with this disease in the OMIM database (which contains a total of 2,387 proteins in the current version). Of the 131 proteins associated with *p,p'*-DDT in the CTD, 48 had associations with asthma (of a total of 1,977 proteins potentially linked to asthma among the 10,509 proteins included in the CTD). The asthma linkage was found only with regard to *p,p'*-DDT; no relationships were found for *p,p'*-DDE, and the single link to *o,p'*-DDT had a nonsignificant *p*-value.

Several links to reproductive phenotypes were retrieved. Female infertility disorders representing diminished ability or inability of a woman to achieve conception, although not further specified in the resources used, were connected with the three chemicals. In parallel, male reproductive disorders, including hypospadias and cryptorchidism, were predicted for *p,p'*-DDT, *o,p'*-DDT, and *p,p'*-DDE. Genes linked to these male phenotypes included the androgen receptor listed for hypospadias. Although more genes were identified from the CTD, the statistical significance decreased. For cryptorchidism, both the androgen receptor and gonadotropin-releasing hormone 1 were predicted. The identities of some of the genes [for details, see Supplemental Material, Table 2 (<http://dx.doi.org/10.1289/ehp.1103533>)] suggested that the DDTs may also have other endocrine-disrupting effects in humans. Type 2 diabetes mellitus appeared to be connected with all three substances via 9 proteins for *p,p'*-DDE (nonsignificant *p*), 40 proteins for *p,p'*-DDT (*p* < 0.0001), and 25 proteins for *o,p'*-DDT (nonsignificant *p*).

For neurobehavioral diseases, Soto and Weaver syndromes were predicted with low *p*-values via the nuclear receptor binding SET domain protein 1 based on OMIM data for all three substances. When using the CTD, additional learning disorders were identified, including learning disorders (*p* = 0.01 for 7 proteins for *p,p'*-DDE; *p* = 0.013 for 12 proteins for *o,p'*-DDT; and *p* < 0.0001 for 25 proteins for *p,p'*-DDT). Similar results were obtained for memory disorders (*p* = 0.05 for 6 proteins for *p,p'*-DDE; and *p* < 0.0001 for 23 proteins for *p,p'*-DDT). These results again reflect the diversity of information present in the two data sources used. Interestingly, *o,p'*-DDT was the only substance studied that predicted a linkage with autism, via the hepatocyte growth factor receptor.

Various cancers appeared to be connected with all three substances. Breast cancer was predicted for all three DDTs using CTD data

and for both parent DDTs using OMIM data (where the link for *p,p'*-DDT is via a single protein, the hyaluronan-mediated motility receptor, *p* = 0.208). Using the CTD, the link for *p,p'*-DDT involved 22 proteins (*p* < 0.001), and for *p,p'*-DDE, 24 proteins (*p* = 0.021), again reflecting the larger amount of information on these chemicals in the CTD. For information on other cancers, see Supplemental Material, Table 2 (<http://dx.doi.org/10.1289/ehp.1103533>).

Discussion

The discipline of systems chemical biology combines experimental findings with computational models with the aim of understanding the effect of xenobiotics on a biological system. This field of research now allows integration of disparate information sources such as high confidence protein–protein association data, protein–disease annotations, “omics” information, and other biological data from databases to explore hidden and unknown connections. Recent advances in toxicogenomics also contribute information to the impact of small molecules on genes and proteins. Because of these developments, advanced computational systems chemical biology models have been developed to decipher the association between environmental chemicals and diseases (Audouze et al. 2010). Accordingly, the combined sources of information can now be applied in computational

models to predict associations between chemical exposures and human health effects. Although such *in silico* prediction of course cannot be considered a proof of causal links, it nonetheless provides justification for hypothesis generation and contributes to interpreting toxicology information from other sources.

The integration of chemical biology and systems biology in a systems chemical biology approach is unsupervised and is based entirely on known chemical and biological information about the behavior of xenobiotics, their interactions with specific proteins, and the consequences in regard to protein–protein interactions and possible disease pathogenesis in humans. The statistical procedure helps in ranking the interactions, but it may not reflect dose–effect relationships. Thus, the disease annotations identified by this *in silico* approach represent hypothetical causal links that need to be explored and verified in a biological setting, whether *in vitro* or *in vivo*, with the aim of deciphering potential toxicity and modes of action of the chemicals. In addition, the links identified are based on current information, which is incomplete, and the absence of a link therefore does not necessarily mean lack of plausibility regarding a particular adverse effect. This problem is clearly illustrated by comparing the predictions generated by using the two databases, with the greater number of predicted associations found using the CTD, which includes less-certain, indirect evidence.




Figure 3. Disease–chemical associations network. The circles represent diseases, with colors representing phenotype categories: red, reproductive disorders; blue, neurodevelopmental-related diseases; green, cancers; gray, other diseases. Rectangles represent the three chemicals studied. The heavier the weight of the connecting lines, the greater the number of proteins linking a chemical to a disease (determined using the OMIM and CTD databases and ChemProt as resources).

Although the smaller number of associations suggested by the OMIM database may be better documented, this likely represents an underestimation because of incomplete information. Although substantial overlap was detected, neither of these databases can therefore be considered definitive, and they may not become so in the foreseeable future. Thus, all predictions depend on current knowledge, much of which is yet uncertain. And again, dose–response relationships cannot be inferred from the predictions. Although these limitations do not invalidate the results, the caveats are important when interpreting the findings. In particular, linkage to asthma and autism, for example, should not be interpreted as indicating that DDT may be a specific cause of these diseases, which are likely multicausal.

In this study, we chose to use the computational model for a major pesticide to assess the usefulness of the computational approach as a tool to bridge gaps in our understanding of environmentally related disease processes by identifying potential mechanistic links. The main results are in close accordance with toxicology findings and results from prospective epidemiological studies that relied on serum concentrations at relevant time windows.

In regard to asthma, the two data sources agree to a substantial extent that the disease may be uniquely associated with the parent DDT compounds. Only nonallergic disease in adult farmers has been linked to DDT usage (Hoppin et al. 2009). In children at 6 years of age, the presence of asthma—independent of atopy—was linked to increased concentrations of *p,p'*-DDE found in umbilical cord serum (Sunyer et al. 2006). Our findings suggest that the epidemiological linkage to *p,p'*-DDE may be indirect and that future studies should also assess exposures to the parent compound.

Antiandrogenic effects of *p,p'*-DDT and *p,p'*-DDE have been demonstrated experimentally (Gray et al. 2001). One epidemiological study showed no association between the anogenital distance in boys and *p,p'*-DDT and *p,p'*-DDE concentrations in maternal serum (Longnecker et al. 2007), whereas another study found a significant association for the latter (Torres-Sanchez et al. 2008). It is not clear from this evidence whether *o,p'*-DDT plays any role in this respect. Most studies on cryptorchidism and hypospadias had limited statistical power or focused only on *p,p'*-DDE (Brucker-Davis et al. 2008). In regard to semen quality, a cross-sectional study of pesticide sprayers currently using DDT showed inverse associations with the current serum concentration, especially for *p,p'*-DDT (Aneck-Hahn et al. 2007), whereas another study involving infertile men showed that the sum of all *p,p'* isomers was negatively associated with sperm concentration (Messaros et al. 2009). As for antiandrogenic

effects studies, much additional information is available on *p,p'*-DDE. Likewise, in regard to type 2 diabetes, epidemiological studies have generally focused on *p,p'*-DDE (ATSDR 2002), but our data suggest that *p,p'*-DDT may be a more likely etiologic agent.

Experimental animal studies document that *p,p'*-DDT is a neurotoxicant, but evidence on other DDT isomers and metabolites is less extensive (ATSDR 2002). A prospective human study in California suggested that the maternal serum concentration of *p,p'*-DDT during pregnancy was a stronger predictor of the neurodevelopment of the child up to 12 months of age than were the *p,p'*-DDE and *o,p'*-DDT concentrations (Eskensazi et al. 2006). Similarly, a study in Spain showed that neuropsychological performance at 4 years of age decreased in children with higher *p,p'*-DDT concentrations in umbilical cord serum (Ribas-Fito et al. 2006). Other studies relied solely on *p,p'*-DDE concentrations (Darvill et al. 2000; Gladen and Rogan 1991; Rogan and Gladen 1991; Sagiv et al. 2008) and may therefore have missed effects associated with the parent compound.

There is sufficient evidence for carcinogenicity of DDT in animals (ATSDR 2002). The main cancer form studied in regard to human DDT exposure is breast cancer. Perhaps the strongest evidence comes from the prospective follow-up of women who provided a blood sample in connection with the Child Health and Development Studies in California in 1959–1967, where 129 women subsequently developed breast cancer before 50 years of age. The odds ratios showed a significant association with *p,p'*-DDT, but not with *o,p'*-DDT or *p,p'*-DDE (Cohn et al. 2007). Support for this notion comes from a Danish study that relied on a 17-year follow-up from 1976: An increased odds ratio for breast cancer was found among women with the highest quartile of serum *p,p'*-DDT concentrations, whereas this tendency was not seen for *p,p'*-DDE (Hoyer et al. 1998). Unfortunately, the majority of studies in this field have relied on DDE measurements, often in cross-sectional designs. Evidence for liver cancer (McGlynn et al. 2006) and testicular cancer (Cohn et al. 2010; Purdue et al. 2009) also supports the notion that *p,p'*-DDT may be the major carcinogen, as associations with *p,p'*-DDE could be due to breakdown of the parent compound. For example, in a case-control study of non-Hodgkin lymphoma using concurrent serum samples, a significant association was seen with *p,p'*-DDE, but the high frequency of nondetectable *p,p'*-DDT concentrations did not allow a proper comparison (Spinelli et al. 2007).

In light of the toxicological and epidemiological evidence on adverse health effects of DDT compounds, this study shows that

the *in silico* approach is highly relevant and meaningful. That said, a major problem in the epidemiological literature is that it mainly links serum concentrations of *p,p'*-DDE to suspected adverse effects. As some of these effects may be due rather to *p,p'*-DDT, the studies therefore rely on a proxy variable for past DDT exposure. However, one cannot assume that all *p,p'*-DDE originates from the subject's own breakdown of the parent compound, and the serum-DDE concentration is therefore imprecise and may be biased. The degree of imprecision will likely vary with the age and time of exposure. Such exposure misclassification generally leads to an underestimation of the true effect of the substances studied. If the effects are ascribed to the unmeasured parent compound, toxicokinetic calculations may perhaps be applied to generate a more appropriate exposure measure that reflects the exposure to the active substance. The opposite error may play a role in toxicological studies, where effects have been attributed to the parent compound, although potentially mediated through a metabolite. However, our results suggest that this possibility is of little significance in regard to DDT.

Overall, our findings demonstrate that the systems chemical biology approach is feasible and may have a pivotal role in considering potential causal associations derived from toxicology and epidemiology studies. Although our approach is based on the current knowledge base and may therefore have overlooked some linkages, the results show that the DDT compounds examined, while chemically related, have tertiary structures, gene expression profiling, and binding properties that deviate sufficiently from one another to predict outcomes that differ substantially. The differences in predicted outcomes are not likely to be due to differences in the amount of information available. Thus, the major parent compound, *p,p'*-DDT, would seem to be much more potent in regard to adverse effects than its isomers and metabolites. In addition, we have identified several new potential target diseases not hitherto examined as relevant outcomes [see Supplemental Material, Table 2 (<http://dx.doi.org/10.1289/ehp.1103533>)]. These potential targets deserve attention in future experimental and epidemiological studies to provide a more complete basis for risk assessment.

The usefulness and validity of the computational approach is likely to improve as more information becomes available, including more chemical–protein data as well as data from “omics” and gene–environment interaction studies. Furthermore, the results of the disease–chemical association analysis will improve in the future as newer, more complete, and curated data become available to expand and fine-tune our understanding of protein–disease associations. In addition, studies like this one will contribute to the

necessary validation of *in silico* approaches and findings, and cumulated experience will help in interpreting such analyses in light of possible unknown interactions and absent dose–effect relationships. Thus, the results obtained with the DDT compounds serve as an illustration of the potential use of computational predictions in toxicology, epidemiology, and environmental health research. The visions expressed by the National Research Council committee on transforming toxicology therefore seem reasonable and realistic.

REFERENCES

- Alibes A, Yankilevich P, Canada A, Diaz-Uriarte R. 2007. IDconverter and IDClight: conversion and annotation of gene and protein IDs. *BMC Bioinformatics* 8:9; doi:10.1186/1471-2105-8-9 [Online 10 January 2007].
- Aneck-Hahn NH, Schulten GW, Bornman MS, Farias P, de Jager C. 2007. Impaired semen quality associated with environmental DDT exposure in young men living in a malaria area in the Limpopo Province, South Africa. *J Androl* 28(3):423–434.
- ATSDR (Agency for Toxic Substances and Disease Registry). 2002. Toxicological profile for DDT, DDE, and DDD. Atlanta, GA:ATSDR.
- Audouze K, Juncker AS, Roque FJ, Krysiak-Baltyn K, Weinhold N, Taboureaux O, et al. 2010. Deciphering diseases and biological targets for environmental chemicals using toxicogenomics networks. *PLoS Comput Biol* 6(5):e1000788; doi:10.1371/journal.pcbi.1000788 [Online 20 May 2010].
- Bader GD, Betel D, Hogue CW. 2003. BIND: the Biomolecular Interaction Network Database. *Nucleic Acids Res* 31(1):248–250.
- Brucker-Davis F, Wagner-Mahler K, Delattre I, Ducot B, Ferrari P, Bongain A, et al. 2008. Cryptorchidism at birth in Nice area (France) is associated with higher prenatal exposure to PCBs and DDE, as assessed by colostrum concentrations. *Hum Reprod* 23(8):1708–1718.
- Cassano A, Manganaro A, Martin T, Young D, Piclin N, Pintore M, et al. 2010. CAESAR models for developmental toxicity. *Chem Cent J* 4(suppl 1):S4.
- Center for Biological Sequence Analysis. 2011. ChemProt Server, version 1. Available: <http://www.cbs.dtu.dk/services/ChemProt/> [accessed 20 September 2010].
- Cohn BA, Cirillo PM, Christianson RE. 2010. Prenatal DDT exposure and testicular cancer: a nested case-control study. *Arch Environ Occup Health* 65(3):127–134.
- Cohn BA, Wolff MS, Cirillo PM, Sholtz RI. 2007. DDT and breast cancer in young women: new data on the significance of age at exposure. *Environ Health Perspect* 115:1406–1414.
- Comparative Toxicogenomics Database (CTD). 2011. Available: <http://ctd.mdibl.org/> [accessed 27 September 2010].
- Darvill T, Lonky E, Reihman J, Stewart P, Pagano J. 2000. Prenatal exposure to PCBs and infant performance on the Fagan Test of Infant Intelligence. *Neurotoxicology* 21(6):1029–1038.
- Davis AP, King BL, Mockus S, Murphy CG, Saraceni-Richards C, Rosenstein M, et al. 2011. The Comparative Toxicogenomics Database: update 2011. *Nucleic Acids Res* 39(Database issue):D1067–D1072.
- de Matos P, Alcantara R, Dekker A, Ennis M, Hastings J, Haug K, et al. 2010. Chemical Entities of Biological Interest: an update. *Nucleic Acids Res* 38(Database issue):D249–D254.
- Eskenza B, Marks AR, Bradman A, Fenster L, Johnson C, Barr DB, et al. 2006. In utero exposure to dichlorodiphenyl-trichloroethane (DDT) and dichlorodiphenyl-dichloroethylene (DDE) and neurodevelopment among young Mexican American children. *Pediatrics* 118(1):233–241.
- Gladen BC, Rogan WJ. 1991. Effects of perinatal polychlorinated biphenyls and dichlorodiphenyl dichloroethene on later development. *J Pediatr* 119(1 Pt 1):58–63.
- Gohlke JM, Thomas R, Zhang Y, Rosenstein MC, Davis AP, Murphy C, et al. 2009. Genetic and environmental pathways to complex diseases. *BMC Syst Biol* 3:46; doi:10.1186/1752-0509-3-46 [Online 5 May 2009].
- Gray LE, Ostby J, Furr J, Wolf CJ, Lambright C, Parks L, et al. 2001. Effects of environmental antiandrogens on reproductive development in experimental animals. *Hum Reprod Update* 7(3):248–264.
- Guldener U, Munsterkotter M, Oesterheld M, Pagel P, Ruepp A, Mewes HW, et al. 2006. MPact: the MIPS protein interaction resource on yeast. *Nucleic Acids Res* 34(Database issue):D436–D441.
- Hamosh A, Scott AF, Amberger JS, Bocchini CA, McKusick VA. 2005. Online Mendelian Inheritance in Man (OMIM), a knowledgebase of human genes and genetic disorders. *Nucleic Acids Res* 33(Database issue):D514–D517.
- Hermjakob H, Montecchi-Palazzi L, Lewington C, Mudali S, Kerrien S, Orchard S, et al. 2004. IntAct: an open source molecular interaction database. *Nucleic Acids Res* 32(Database issue):D452–D455.
- Hoppin JA, Umbach DM, London SJ, Henneberger PK, Kullman GJ, Coble J, et al. 2009. Pesticide use and adult-onset asthma among male farmers in the Agricultural Health Study. *Eur Respir J* 34(6):1296–1303.
- Hoyer AP, Grandjean P, Jorgensen T, Brock JW, Hartvig HB. 1998. Organochlorine exposure and risk of breast cancer. *Lancet* 352(9143):1816–1820.
- InParanoid: Eukaryotic Ortholog Groups. 2009. InParanoid version 7. Available: <http://inparanoid.sbc.su.se/cgi-bin/index.cgi> [accessed 26 June 2009].
- Joshi-Tope G, Gillespie M, Vastrik I, D'Eustachio P, Schmidt E, de Bono B, et al. 2005. Reactome: a knowledgebase of biological pathways. *Nucleic Acids Res* 33(Database issue):D428–D432.
- Judson R, Richard A, Dix D, Houck K, Elloumi F, Martin M, et al. 2008. ACToR—Aggregated Computational Toxicology Resource. *Toxicol Appl Pharmacol* 233(1):7–13.
- Kanehisa M, Goto S, Hattori M, Aoki-Kinoshita KF, Itoh M, Kawashima S, et al. 2006. From genomics to chemical genomics: new developments in KEGG. *Nucleic Acids Res* 34(Database issue):D354–D357.
- Lage K, Hansen NT, Karlberg EO, Eklund AC, Roque FS, Donahoe PK, et al. 2008. A large-scale analysis of tissue-specific pathology and gene expression of human disease genes and complexes. *Proc Natl Acad Sci USA* 105(52):20870–20875.
- Lage K, Karlberg EO, Stirling ZM, Olason PI, Pedersen AG, Regina O, et al. 2007. A human phenotype-interactome network of protein complexes implicated in genetic disorders. *Nat Biotechnol* 25(3):309–316.
- Lage K, Mollard C, Greenway S, Wakimoto H, Gorham JM, Workman CT, et al. 2010. Dissecting spatio-temporal protein networks driving human heart development and related disorders. *Mol Syst Biol* 6:381; doi:10.1038/msb.2010.36 [Online 22 June 2010].
- Lhasa Limited. 2011. Derek Nexus. Available: https://www.lhasalimited.org/index.php/derek_nexus/ [accessed 29 July 2011].
- Liu T, Lin Y, Wen X, Jorissen RN, Gilson MK. 2007. BindingDB: a web-accessible database of experimentally determined protein-ligand binding affinities. *Nucleic Acids Res* 35(Database issue):D198–D201.
- Longnecker MP, Gladen BC, Cupul-Uicab LA, Romano-Riquer SP, Weber JP, Chapin RE, et al. 2007. In utero exposure to the antiandrogen 1,1-dichloro-2,2-bis(p-chlorophenyl)ethylene (DDE) in relation to anogenital distance in male newborns from Chiapas, Mexico. *Am J Epidemiol* 165(9):1015–1022.
- McGlynn KA, Abnet CC, Zhang M, Sun XD, Fan JH, O'Brien TR, et al. 2006. Serum concentrations of 1,1,1-trichloro-2,2-bis(p-chlorophenyl)ethane (DDT) and 1,1-dichloro-2,2-bis(p-chlorophenyl)ethylene (DDE) and risk of primary liver cancer. *J Natl Cancer Inst* 98(14):1005–1010.
- Messaros BM, Rossano MG, Liu G, Diamond MP, Friderici K, Nummy-Jernigan K, et al. 2009. Negative effects of serum p,p'-DDE on sperm parameters and modification by genetic polymorphisms. *Environ Res* 109(4):457–464.
- Mishra GR, Suresh M, Kumar K, Kannabiran N, Suresh S, Bala P, et al. 2006. Human protein reference database—2006 update. *Nucleic Acids Res* 34(Database issue):D411–D414.
- National Research Council. 2007. Toxicity Testing in the 21st Century: A Vision and a Strategy. Washington, DC:National Academies Press.
- NCBI (National Center for Biotechnology Information). 2011a. EntrezGene. Available: <http://www.ncbi.nlm.nih.gov/gene> [accessed 23 September 2010].
- NCBI (National Center for Biotechnology Information). 2011b. OMIM – Online Mendelian Inheritance in Man. Available: <http://www.ncbi.nlm.nih.gov/omim> [accessed 29 September 2010].
- O'Brien KP, Remm M, Sonnenhammer EL. 2005. InParanoid: a comprehensive database of eukaryotic orthologs. *Nucleic Acids Res* 33(Database issue):D476–D480.
- Oprea TI, Tropsha A, Faulon JL, Rintoul MD. 2007. Systems chemical biology. *Nat Chem Biol* 3(8):447–450.
- Pagel P, Kovac S, Oesterheld M, Brauner B, Dunger-Kaltenbach I, Frishman G, et al. 2005. The MIPS mammalian protein–protein interaction database. *Bioinformatics* 21(6):832–834.
- Pavan M, Worth AP. 2008. Publicly-accessible QSAR software tools developed by the Joint Research Centre. *SAR QSAR Environ Res* 19(7–8):785–799.
- Purdue MP, Engel LS, Langseth H, Needham LL, Andersen A, Barr DB, et al. 2009. Prediagnostic serum concentrations of organochlorine compounds and risk of testicular germ cell tumors. *Environ Health Perspect* 117:1514–1519.
- R Development Core Team. 2010. R: A Language and Environment for Statistical Computing. Vienna:R Foundation for Statistical Computing, version R-2.11.0. Available: <http://www.r-project.org> [accessed 22 April 2010].
- Ribas-Fito N, Torrent M, Carrizo D, Munoz-Ortiz L, Julvez J, Grimalt JO, et al. 2006. In utero exposure to background concentrations of DDT and cognitive functioning among preschoolers. *Am J Epidemiol* 164(10):955–962.
- Rogan WJ, Gladen BC. 1991. PCBs, DDE, and child development at 18 and 24 months. *Ann Epidemiol* 1(5):407–413.
- Roth BL, Lopez E, Beischel S, Westkaemper RB, Evans JM. 2004. Screening the receptorome to discover the molecular targets for plant-derived psychoactive compounds: a novel approach for CNS drug discovery. *Pharmacol Ther* 102(2):99–110.
- Rual JF, Venkatesan K, Hao T, Hirozane-Kishikawa T, Dricot A, Li N, et al. 2005. Towards a proteome-scale map of the human protein–protein interaction network. *Nature* 437(7062):1173–1178.
- Sagiv SK, Nugent JK, Brazelton TB, Choi AL, Tolbert PE, Altschul LM, et al. 2008. Prenatal organochlorine exposure and measures of behavior in infancy using the Neonatal Behavioral Assessment Scale (NBAS). *Environ Health Perspect* 116:666–673.
- Salwinski L, Miller CS, Smith AJ, Pettit FK, Bowie JU, Eisenberg D. 2004. The Database of Interacting Proteins: 2004 update. *Nucleic Acids Res* 32(Database issue):D449–D451.
- Spinelli JJ, Ng CH, Weber JP, Connors JM, Gascoyne RD, Lai AS, et al. 2007. Organochlorines and risk of non-Hodgkin lymphoma. *Int J Cancer* 121(12):2767–2775.
- Stark C, Breitkreutz BJ, Reguly T, Boucher L, Breitkreutz A, Tyers M. 2006. BioGRID: a general repository for interaction datasets. *Nucleic Acids Res* 34(Database issue):D535–D539.
- Sunyer J, Torrent M, Garcia-Esteban R, Ribas-Fito N, Carrizo D, Romieu I, et al. 2006. Early exposure to dichlorodiphenyl-dichloroethylene, breastfeeding and asthma at age six. *Clin Exp Allergy* 36(10):1236–1241.
- Taboureaux O, Nielsen SK, Audouze K, Weinhold N, Edsgard D, Roque FS, et al. 2011. ChemProt: a disease chemical biology database. *Nucleic Acids Res* 39(Database issue):D367–D372.
- Torres-Sanchez L, Zepeda M, Cebrán ME, Belkind-Gerson J, García-Hernández RM, Belkind-Valdivinos V, et al. 2008. Dichlorodiphenyl-dichloroethylene exposure during the first trimester of pregnancy alters the anal position in male infants. *Ann N Y Acad Sci* 1140:155–162.
- Wheeler DL, Barrett T, Benson DA, Bryant SH, Canese K, Chetvernin V, et al. 2007. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* 35(Database issue):D5–D12.
- Zanzoni A, Montecchi-Palazzi L, Quondam M, Ausiello G, Helmer-Citterich M, Cesareni G. 2002. MINT: a Molecular INteraction database. *FEBS Lett* 513(1):135–140.